

[Crichton Browne J \(1872\)](#). Conium in the treatment of acute mania. *Lancet* 1:217-8.

Key passages

CONIUM IN THE TREATMENT OF ACUTE MANIA.

BY J. CRICHTON BROWNE, M.D., F.R.S.E.,
MEDICAL DIRECTOR, WEST RIDING ASYLUM.

(Concluded from page 183.)

WITH the view of determining the relative duration of cases of acute mania treated with conium and with other remedies, I have contrasted twelve cases from each category. As the result of that contrast, I find that twelve consecutive cases treated from the first with conium, admitted subsequent to Feb. 25th, 1871, and since discharged recovered, had an average duration, as measured by residence in the asylum, of 102 days; whereas twelve consecutive cases treated with other remedies, such as bromide of potassium, cannabis indica, chloral, and digitalis, admitted subsequent to October 1st, 1870, and since discharged recovered, had an average duration, as measured by residence in the asylum, of 150 days. This shows a balance of 48 days in favour of conium. It will be seen, from the cases reported below, that when conium was used, that period of convalescence which it is thought prudent to interpose between the date when recovery might be called complete and the date of discharge from the asylum was in no instance unusually shortened. In some cases it might be thought that it was unnecessarily prolonged. So rapid and decisive were the beneficial effects of the remedy that it was feared at first they might prove transitory, and that, as is not seldom the case when recovery is sudden, a relapse might be looked for. Experience having now taught that this fear is groundless, a considerable curtailment of the term of medical supervision may be henceforth safely conceded. Indeed, so prompt is recovery, as a rule, under the conium treatment, that it appears that the necessity of removal to an asylum may be obviated in some cases of acute mania if it is had recourse to, and judiciously conducted, in their initial stage.