

Marks HM. The Kendrick-Eldering-(Frost) pertussis vaccine field trial

Commentary on: Kendrick P, Eldering G (1939). A study in active immunization against pertussis. American Journal of Hygiene 29: 133-153.

Cite as: Marks HM (2006). The Kendrick-Eldering-(Frost) pertussis vaccine field trial. The James Lind Library (www.jameslindlibrary.org). Accessed Wednesday 9 May 2007.

©Harry Marks, 2006

Author contact details: Harry M Marks, History of Medicine, The Johns Hopkins University, 1900 E. Monument St, Baltimore, MD 21205, USA. Email: hmarks@jhmi.edu

In 1939, Pearl Kendrick and Grace Eldering reported the results of a field trial of a new pertussis vaccine that the two women had developed at the Michigan Department of Health ([Kendrick and Eldering 1939](#)). The trial had run for more than three years (March 1934 - November 1937), enrolling 5,815 children. The vaccinated group "was made up of all children of acceptable age and history who presented themselves at the city immunization clinics for pertussis vaccination." The control group "was selected at random" from a list of non-immunized children maintained by the Grand Rapids City Health Department, producing an "approximately equal sample of children of the same age, and in the same districts as the injected children" (Kendrick and Eldering 1939: 133-136). Based on a careful and innovative analysis of events/person-years at risk, the authors concluded that "substantial protection had been afforded to the injected group":

"...there were 2.3 annual attacks per 100 [person-years] in the injected group and 15.1 in the control group. Expressed statistically, the difference is equal to fifteen times the standard error, indicating that the probability of such a difference occurring by chance alone is one in several million trials" (Kendrick and Eldering 1939: 152-153).

Methodologically, the original field trial design was flawed. The experimental group was self-selected and only control subjects were randomly chosen. Despite careful attention paid to case detection and diagnosis, 1603 observations from the study's early years had to be excluded from the final analysis. Several features of the trial nonetheless make it an important contribution, not simply to the development of an effective pertussis vaccine, but to the history of controlled trials:

1. The trial was enabled at the height of the Great Depression by a diverse coalition of governmental (local, state and federal) and private agencies, whose activities were orchestrated by Kendrick and Eldering (Shapiro-Shapin 2007);
2. The trial was unusual for the level of attention given to case diagnosis and follow-up, and to the discussion of unknown factors which might have biased the results;
3. A similar level of detail was given in reporting on the analysis and the methodological limitations of the field trial;
4. It introduced to clinical trials a methodology (events/time analysis) more commonly found in observational epidemiology;
5. The trial provided an influential network of specialists in public health and biostatistics an occasion to articulate their views of what constituted a "well-controlled" trial.

Pertussis vaccines: a brief history

In 1913, Charles Nicolle introduced the first in a series of pertussis vaccines developed over the next two decades (Nicolle and Connor 1913). Few of these vaccines produced consistent results, leading the Council on Pharmacy and Chemistry of the American Medical Association to withdraw its approval from all commercially produced vaccines in 1931 (Council on Pharmacy and Chemistry 1931). Pertussis was then increasingly a dominant cause of childhood mortality, outranking diphtheria, scarlet fever and measles in many European countries (League of Nations 1934). In the late 1920s, Danish researchers, armed with a better understanding of the antigenic

properties of different pertussis strains, had introduced a new generation of vaccines. These European reports of favorable results sparked renewed North American interest in vaccine development: Lewis Sauer, Pearl Kendrick and James Doull were among the researchers who began experimenting with new techniques for developing a more potent, clinically consistent pertussis vaccine (McDonald 1935). Vaccine evaluation had not followed suit. Existing studies had "been carried on in the medical literature with slight regard for properly controlled experimentation" (Lawson 1932: 4-5).

The investigators, the community and the trial

Born in 1890, Pearl Kendrick spent several years after college teaching high school science in New York State. She worked briefly at the New York State Department of Health before moving to the Michigan department in 1920, where she worked with Reuben Kahn in developing a new diagnostic assay for syphilis. Six years later, she took charge of the Department's Western Michigan Laboratory. In 1932, she received a ScD in bacteriology from the Johns Hopkins School of Hygiene and Public Health (Cattell and Cattell 1938; Allen 1979: 12). On her return to Michigan she asked CC Young, state Director of Laboratories, for permission to work on "whooping cough"; he allegedly replied "If you are having any fun working on whooping cough, go ahead" (Eldering 1971:200). Accompanied by bacteriologist Grace Eldering, who had joined the Department's laboratories in 1928, Kendrick "set out" every day after "the laboratory closed" to collect *b. pertussis* specimens. As Eldering later put it:

"We learned about the disease and the depression at the same time. Many of the families were very poor and their living conditions pitiful. [...] We listened to sad stories told by desperate fathers who could find no work. We collected specimens by the light of kerosene lamps, from whooping, vomiting, strangling children. We saw what the disease could do. In the laboratory we isolated the pertussis bacillus, not from every patient, but from most of them in the early stages of the disease. [...] The cultures were saved and studied in every possible way" (Eldering 1971:200).

The Michigan department was one of the few public health laboratories in the United States with a program in vaccine development and production.ⁱ With CC Young's encouragement, Kendrick and Eldering began working on a pertussis vaccine. The two bacteriologists had closely followed the new literature concerning the importance of using recent bacterial cultures to cultivate pertussis vaccine. By January 1933, they had prepared their first samples (Eldering 1971:201);ⁱⁱ by mid-year they were drawing up plans for an elaborate field trial in Grand Rapids, Michigan.

The Great Depression was perhaps the worst time in recent history to start a large, complex clinical trial: local government funding for public health, never magnanimous, was sharply curtailed (Allen 1979:14). Kendrick cobbled together financial support from federal emergency relief programs, state and city health departments, city government and private donors. Official funds were supplemented with volunteer labor (nurses, technicians, and private physicians) who worked well beyond a paid forty-hour week in preparing, testing, and administering the vaccine. Even so, periodic funding crises hampered case follow-up and record keeping for the study (Eldering 1971).ⁱⁱⁱ

Trained as laboratory researchers in bacteriology, neither Kendrick nor Eldering had any prior experience with clinical trials. The trial's design and procedures were a work-in-progress, modified as operational and analytical difficulties became apparent. Crucial to all versions of the study was a federally funded 1934 "pre-school immunization survey" for Grand Rapids, which identified the exposure history of the city's under-five-year-old population. The initial plan called for vaccine to be given to a test group of "non-immune" children "exposed in their own home," with controls ideally to be chosen from other children in the same family. When it proved impossible to reach "exposed" children with the vaccine before they became symptomatic, a more elaborate procedure was adopted.

The experimental group consisted of children who voluntarily came to the city's health clinics to be inoculated. The control group was drawn from non-immune children "from the same district" in the city as the vaccinated child.^{iv} Nurses from the Grand Rapids Health Department assigned to one or more of its 18 health districts did the follow-up to identify cases of whooping cough among study participants (Eldering 1971; Kendrick and Eldering 1939). Initially, the nurses also selected children for the control group; this task was later transferred to Kendrick's office, which relied on the immunization survey. The initial emphasis on selecting controls from the families of inoculants was similarly dropped over time.^v In October 1935, Kendrick and Eldering presented their preliminary results at the annual American Public Health Association meeting. Of 1592 children enrolled at that time (712 in "test group"; 880 controls), there were 63 cases of whooping cough in the control group, and only 3 in the vaccinated group (Kendrick and Eldering 1936).

Such favorable results contrasted strongly with the experiences of other American researchers, especially a contemporaneous Cleveland study being led by James Doull. Doull and his colleagues (1936) were conducting a controlled study of their pertussis vaccine, enrolling children 6-15 months old who visited the city's free milk distribution centers. Assignment was by alternation: the first eligible child was offered vaccination, the second

served as a control. Because of substantial losses to enrollment (refusals, drop-outs, and doubtful pre-trial histories of exposure), they soon switched to a 2:1 assignment ratio, with every third eligible child serving as a control. After 979 children had been enrolled, they were unable to detect any advantage for the vaccine: 61 cases/483 children (test group) versus 71 cases/496 children (247 controls plus 249 refusals) ([Doull et al. 1936](#)).

Doull had considerable standing in the public health research community: he was Professor of Hygiene at Western Reserve University, a former Associate Professor of Epidemiology at the Johns Hopkins School of Hygiene, and first author of a randomized trial assessing the effects of irradiation with ultraviolet light on the incidence of the common cold ([Doull 1931](#)). Against a background of skepticism generated by disappointments with earlier pertussis vaccines, the differences between Doull's preliminary findings and Kendrick's results presented a problem for public health authorities. Among the differences between the two studies, children in Doull's trial were markedly younger (<15 months old), he relied more heavily on physicians for diagnosis of cases, and fewer (435) children were enrolled. Importantly, Doull selected vaccine and control children from the same pool - infants attending Cleveland's milk distribution stations ([Doull et al. 1936](#)).

Senior figures at the American Public Health Association (APHA) Committee on Administrative Practices (headed by Haven Emerson, former New York City health commissioner and dean of the Columbia School of Public Health) soon found themselves involved in complex negotiations to assess the strengths (and weaknesses) of Kendrick's ongoing field trials. The Committee, engaged in evaluating a variety of existing public health practices, did not want to endorse an inadequately tested vaccine.

The consultant

Although Doull did not present his findings publicly until the 1936 meeting of the APHA, he was already in touch with Kendrick about their conflicting findings.^{vi} Early in 1936, they had discussed the possibility of meeting with Wade Hampton Frost, professor of epidemiology at Johns Hopkins, to review their respective studies. His personal status as America's premier epidemiologist aside, Frost was not an obvious choice to advise on vaccine field trials: he was known mainly for his pioneering work in observational studies of polio, influenza and tuberculosis (Daniel 2004). Frost's other main activities included the development of the Eastern Health District and Hagerstown community studies, and work on the mathematical modeling of epidemics, done in collaboration with statistician Lowell Reed ("Introduction" 1941; Fee 1987).

Kendrick was nonetheless enthusiastic about getting Frost's advice, help that Frost was not eager to provide. Whether because of the press of other work, because of his close relationship with Doull, a former student and colleague, or for some other reason, Frost was reluctant to get involved. Thus, in a letter to Kendrick, he wrote:

"I cannot think that any judgment other than Dr. Doull's or your own is really needed for interpretation of your data, and on general principles, I think those who know the data most intimately are best qualified to interpret them."^{vii}

APHA leaders appealed to Frost to intervene: in addition to Haven Emerson, these included George Ramsey, head of communicable diseases at the New York State Department of Health, previously at the Michigan department and, like Doull, a former student and colleague of Frost's. The APHA committee overseeing immunization practices was deadlocked. According to Ramsey, Doull himself "is convinced that the Cleveland study has been so carefully and well done that the matter is settled."^{viii}

Over the summer and early fall, Frost was reluctantly drawn into discussions about Kendrick's trial. Still, he continued to hope that the issue could be resolved without a site visit:

"I very strongly suspect that Miss Kendrick's field studies are not set up in such a way as to give a really good control. [...] Not 1 out of 10 - perhaps not one out of 50 - attempts is successful and as a mere matter of probability the odds are strongly against Miss Kendrick's experiment being sound. If this is the case it may be that her own account of her procedure, which I hope to have from her in writing, will show some basic defect. In that case, it will not be necessary for anyone to go to Grand Rapids."^{ix}

In the event, Frost went to Grand Rapids twice (in November 1936 and September 1937), and offered a detailed critique of the trial and a plan for analyzing the results. Frost identified several key issues:^x

1. Owing to the long, slow buildup of the trial, the study population overall was quite heterogeneous. Frost recommended dropping the children vaccinated by private physicians, whose records were less reliable than those from the city clinics.
2. Similarly, in the early years of the trial, follow-up of control children was either inadequate or the records were incomplete. Frost ultimately recommended that the first 1100 observations be dropped from the

analysis.

3. Because recruitment to the trial varied over the life of the study, as did the frequency of nursing visits to look for whooping cough, Frost suggested examining the data using an innovative event/time analysis, which would calculate the number of months each child (vaccinated or control) was at risk for developing pertussis. In keeping with the event analysis, Frost proposed that a child might be a control subject one month and an experimental subject the following month, if they happened to receive the vaccine in the interim.
4. The possibility of unknown differences between experimental and control groups, because of differences in the way they had been recruited, continued to trouble Frost. He pushed hard for any analyses which would comfort a suspicious mind like his. For example, were rates of other communicable diseases also lower in the experimental group (as might be expected if the vaccinated children were from a higher socio-economic group than the controls)? Kendrick made the recommended comparison of non-pertussis disease incidence, but found no differences between the groups in levels of other childhood infectious diseases (Kendrick and Eldering 1939:142).

Nor was Frost alone in his concerns about the trial's design. Harvard statistician Edwin Wilson, drawn into the discussion by George Lawson, a Harvard alumnus involved in pertussis research, sympathized with Frost's reservations. As Wilson commented, there were some observable differences between the experimental and control groups. They might be important but then again they might not be:

"...one can never be certain but what such a difference has some significance with respect to some hidden factor which makes the controls not a fair control. [...] I can see how a cautious person like Frost might think that you weren't getting enough to pay to go on." ^x

Whatever his initial reservations, it is clear from the tone of Frost's correspondence that his professional regard for Kendrick's commitment to analytical rigor increased considerably over time, as did his regard for the Michigan data. ^{xii} As he wrote Haven Emerson:

"...considering the circumstances under which the work had to be done - beginning on a small scale and expanding as new funds permitted - I do not see how the complications could well have been avoided. The effect of all these circumstances is that eventual analysis of the material must be considerably more complicated than is required, for instance, in Dr. Doull's much simpler material. However, the records are such as to permit such analysis as may be required and, though the statistical work involved will be somewhat laborious, I see no reason why the end result should not be definite." ^{xiii}

In consultation with Kendrick, Eldering and a medically qualified epidemiologist, FC Forsbeck, Frost worked out a detailed plan for analysing the data. ^{xiv} The ensuing statistical work was laborious; the investigators produced a 30 + page codebook detailing the handling of each record. After deciding which records could be used, the investigators had then to transfer the data for 5815 patients, month by month, to punch cards in order to accomplish Frost's event-time analysis. ^{xv}

Frost was among the many individuals, outside and inside the Michigan department, who received a draft of the paper for comment. ^{xvi} The published paper followed meticulously Frost's memoranda about analysis and reporting of the trial data, and his guidance was graciously acknowledged (although one could not possibly guess the extent of his contributions from this note). By the time the report was published in May 1939, Frost, its invisible co-author, had died, the victim of an esophageal cancer that had appeared after his September 1937 visit to Michigan.

Publication of Kendrick and Eldering's paper did not resolve questions about the value of pertussis vaccine. There was general agreement in the specialist community that a replication of Kendrick's study was desirable. Kendrick, in collaboration with James Doull, worked out a proposal for a new field study, which APHA leaders proved reluctant to fund, preferring to wait on corroboration from other, smaller community studies. ^{xvii} The technical design of pertussis vaccines was, in any case in rapid evolution, and it was left to Kendrick and other researchers to conduct field trials of newer vaccines in the 1940s and 1950s ([Kendrick 1942](#); [Kendrick 1943](#); [Bell 1948a](#); [Bell 1948b](#); [MRC 1951](#); [Jefferson 2006](#)).

The individuals involved in the pertussis field trials of the 1930s - Pearl Kendrick, Grace Eldering, Wade Hampton Frost, James Doull et al. - do not figure in any of the standard histories of clinical trials (Bull 1959; Lilienfeld 1982; Matthews 1995; Marks 1997; Chalmers 2001). Yet Kendrick's skill in assembling the people and money to conduct a large field trial in the midst of the Great Depression (Shapiro-Shapin 2007) is as remarkable as her dedication to finding out whether her vaccine really did work. In addition to her response to Frost's criticisms,

time and again, Kendrick welcomed close scrutiny of her data.^{xviii} The 1934 pertussis vaccine trial launched her national career in vaccine development and standardization, which continued for many decades (Eldering 1971; Brown 1971). Among her many accomplishments was the development of the mouse protection test, which had a long international history as a measure of vaccine potency (Kendrick et al 1949; Murata 1971; Jefferson 2006).

The most surprising character to figure in this story remains Wade Hampton Frost. Unlike other epidemiologists and biostatisticians of his generation - Edgar Sydenstricker, Edwin Wilson, Major Greenwood - Frost never wrote a line about experimentation, controlled or otherwise. Yet it is clear from the record of his encounters with Pearl Kendrick that he had a profound appreciation of the importance of controlled comparison in any epidemiological analysis. And he offered to Kendrick and Eldering the methodological fruits of his epidemiological studies of tuberculosis, influenza and polio. For Frost, in any epidemiological analysis, it was crucial to reliably capture the period over which events occurred (Frost 1936). While Frost did not invent the methodology of "event/person-years" analysis (Vandenbroucke 1985), he had a profound appreciation for defining the period at risk in any epidemiological analysis, including Kendrick's field trial of pertussis vaccines.

Several years after Kendrick and Eldering's field trial, Joseph Bell, another epidemiologist trained at Johns Hopkins, published a field trial of a new pertussis vaccine developed by the US Public Health Service. Bell's study avoided many of the pitfalls in design which had hampered earlier field trials, giving careful attention to random selection of experimental and control groups and to avoiding the effects of ascertainment bias in follow-up of results (Bell 1941; Chalmers 2006). It is not possible to establish whether Bell had direct contact with Frost during his initial time at Johns Hopkins. We do know that Frost's preferred method of teaching was the "laboratory problem", in which students would work through the planning and analysis of an epidemiological study, discussing methodological issues as they proceeded ("Introduction" 1941). Whether directly or indirectly through one of Frost's colleagues, it seems likely that the lessons of Frost's experience with Kendrick's field trial were passed on.

Neither Frost nor Johns Hopkins appear in the standard histories of clinical trials, most of which refer to RA Fisher and Austin Bradford Hill, British statisticians, the British Medical Research Council, and/or the long tradition - as illustrated by the records published in the James Lind Library - of medical investigators seeking 'fair' and objective tests of therapies. Yet in retrospect, it is noticeable how many innovative studies of the 1930s and 1940s were conducted by individuals with a strong connection to Wade Frost and/or to his Johns Hopkins colleague, the statistician Lowell Reed. In 1931, James Doull had conducted a randomized trial to assess whether ultraviolet light would reduce the incidence of common colds, using, on Reed's advice, an innovative urn device to assign patients to experimental and control groups (Doull 1931). Carroll Palmer guided the US Public Health Service randomized trial of streptomycin (Long and Ferebee 1950), which was conducted contemporaneously with the more famous MRC study (MRC 1948). And Margaret Merrell was statistician for the wartime penicillin clinical evaluations and a prominent postwar advocate and popularizer of controlled clinical trials in North America (Marks 1997). Along with Wade Hampton Frost's role as Pearl Kendrick's mentor in analysing her 1934 field trial of pertussis vaccine, they suggest the existence of a strong and hitherto unacknowledged network in the development of clinical trial methodology and practice.

This James Lind Library commentary has been republished in the *Journal of the Royal Society of Medicine* 2007; 100: 242-247.

Acknowledgements: I am grateful to Iain Chalmers for alerting me to Kendrick's pertussis trials and for comments on earlier drafts; to Kay Dickersin, Jan Vandenbroucke, and Tom Jefferson for advice, and to archivists too numerous to mention.

References

A. Manuscript Sources

Grace Eldering papers, Library of Michigan, Lansing, MI.

Pearl Kendrick papers, Bentley Historical Library, University of Michigan, Ann Arbor, MI

Michigan Department of Public Health, Bureau of Laboratories, Record Group 76-73, Library of Michigan, Lansing, MI.

Edwin Bidwell Wilson papers [GA94.10], Harvard Medical Library in the Francis A. Countway Library of Medicine, Boston, MA.

Charles-Edward Amory Winslow papers, Sterling Memorial Library, Yale University, New Haven, CT.

B. Published Sources

- Allen MTS (1979). Bureau of Laboratories. Michigan Department of Health: History 1907-1975. Michigan Department of Health. Copy on deposit at the Bentley Historical Library, University of Michigan, Ann Arbor, MI.
- Bell JA (1941). Pertussis prophylaxis with two doses of alum-precipitated vaccine. *Public Health Reports* 56: 1535-1546.
- Bell JA (1948a). Use of an alum-precipitated mixture of pertussis vaccine and diphtheria toxoid. *JAMA* 137: 1009-1016.
- Bell JA (1948b). Use of two doses of an alum-precipitated mixture of diphtheria toxoid and pertussis vaccine. *JAMA* 137: 1276-1281.
- Brown GC (1971). Symposium on pertussis immunization, in honor of Dr. Pearl L. Kendrick in her eightieth year: Current Status of Combined Vaccines. *Health Laboratory Sciences* 8: 210-214.
- Bull JP (1959). The historical development of clinical therapeutic trials. *Journal of Chronic Diseases* 10: 218-248.
- Cattell JM, Catell J (1938). *American Men of Science*. 6th edition. The Science Press, p 767.
- Chalmers I (2001). Comparing like with like: some historical milestones in the evolution of methods to create unbiased comparison groups in therapeutic experiments. *International Journal of Epidemiology* 30: 1156-1164.
- Chalmers I (2006). Joseph Asbury Bell and the birth of randomized trials. In: *The James Lind library* (www.jameslindlibrary.org).
- Council on Pharmacy and Chemistry, AMA (1931). Pertussis vaccines omitted from NNR. *JAMA* 96: 613.
- Daniel TM (2004). *Wade Hampton Frost. Pioneer Epidemiologist 1880-1938*. University of Rochester Press.
- Doull JA, Hardy M, Clark J, Herman NB (1931). The effect of irradiation with ultra-violet light on the frequency of attacks of upper respiratory diseases (common cold). *American Journal of Hygiene* 13: 460-477.
- Doull JA, Shibley GS, McClelland JE (1936). Active immunization against whooping cough. Interim Report of the Pertussis Experience. *American Journal of Public Health* 26: 1097-1105.
- Eldering G (1971). Symposium on pertussis immunization, in honor of Dr. Pearl L Kendrick in her eightieth year: Historical notes on pertussis immunization. *Health Laboratory Sciences* 8: 200-205.
- Fee E (1987). *Disease & discovery. A History of the Johns Hopkins School of Public Hygiene and Public Health, 1916-1939*. Johns Hopkins University Press.
- Frost WH (1936). How much control of tuberculosis? *American Journal of Public Health* 28: 7-13.
- "Introduction" (1941). In: Maxcy KF, ed. *Papers of Wade Hampton Frost MD*. The Commonwealth Fund, pp 1-20.
- Jefferson T (2006). Why the MRC randomised trials of whooping cough (pertussis) vaccines remained important 40 years after they were done. *The James Lind Library* (www.jameslindlibrary.org).
- Kendrick P (1942). Use of alum-treated pertussis vaccine, and of alum-precipitated combined pertussis vaccine and diphtheria toxoid for active immunization. *American Journal of Public Health* 32: 615-626.
- Kendrick P (1943). A field study of alum-precipitated combined pertussis vaccine and diphtheria toxoid for active immunization. *American Journal of Hygiene* 38: 193-202.
- Kendrick P, Eldering G (1936). Progress report on pertussis immunization. *American Journal of Public Health* 26: 8-12.
- Kendrick P, Eldering G (1939). A study in active immunization against pertussis. *American Journal of Hygiene* 29 Sec. B: 133-153.

Kendrick P, Updyke, EL, Eldering G (1949). Comparison of pertussis cultures by mouse protection and virulence tests. *American Journal of Public Health* 39:179-184.

Lawson GM (1932). Studies on bacillus pertussis. DPh thesis, Harvard School of Public Health, Boston, MA.

League of Nations (1934). Relative importance of the principal infectious diseases of childhood (diphtheria, scarlet fever, whooping-cough and measles. *Epidemiological Report No. 173, Table III.*

Lilienfeld AM (1982). *Ceteris Paribus*: the evolution of the clinical trial. *Bulletin of the History of Medicine* 56:1-18.

Long ER, Ferebee SH (1950). A controlled investigation of streptomycin treatment in tuberculosis. *Public Health Reports* 65:1421-1451.

McDonald FC (1935). Whooping cough, with particular reference to prophylaxis and treatment with vaccines. *New England Journal of Medicine* 213:198-203.

Marks HM (1997). The progress of experiment. Science and therapeutic reform in the United States, 1900-1990. Cambridge University Press.

Matthews JR (1995). Quantification and the quest for medical certainty. Princeton University Press.

Medical Research Council (1948). Streptomycin treatment of pulmonary tuberculosis: a Medical Research Council investigation. *BMJ* 2:769-782.

Medical Research Council (1951). The prevention of whooping-cough by vaccination. *BMJ* 1:1463-1471.

Murata R, Perkins FT, Pittman M, Scheibel I, Sladkey K (1971). International collaborative studies on the pertussis vaccine potency assay. Part played by the challenge in the mouse-protection test. *Bulletin World Health Organization* 44:673-687.

Nicolle C, Conor A (1913). Vaccinothérapie dans le coqueluche. *Comptes rendus de l'Academie des Sciences de Paris* 16:1849-1851.

Shapiro-Shapin CG (2007). A whole community working together: Pearl Kendrick, Grace Eldering and the Grand Rapids Pertussis Trials, 1932-1939. *Michigan Historical Review*, 33 (in press).

Vandenbroucke JP (1985). On the rediscovery of a distinction. *American Journal of Epidemiology* 121:627-628.

Endnotes

- i. On the Michigan laboratory under Young, see Allen (1979), pp. 7-29. Other health departments with active research programs would include Massachusetts, New York State and New York City. See: Rosenkrantz BG (1972). *Public health and the state. Changing views in Massachusetts, 1846-1936.* Harvard University Press and Sexton AM (1967). *A chronicle of the Division of Laboratories & Research.* New York State Department of Health. The Stinehour Press.

[Back](#)

- ii. Pearl Kendrick to Louis Sauer, June 12, 1934. Box 1, Grace Eldering papers.

[Back](#)

- iii. Pearl Kendrick, The Status of the Study of Whooping Cough in Grand Rapids, December 10, 1934; Pearl Kendrick to CC Young, August 27, 1935, both Box 14, Michigan Department of Health; Landmarks in Pertussis Immunization Study, 1932-1935; Pearl Kendrick to Wade Hampton Frost, September 15, 1937, both Box 4, Kendrick papers.

[Back](#)

- iv. Michigan Department of Health, Bureau of Laboratories, Whooping Cough Prevention Study, 1-20-34; idem, Whooping Cough Prevention Study, General Plan, revised 10-16-34; idem, Whooping Cough Prevention

Study, 10-13-1936, all Box 4, Kendrick papers; Pearl Kendrick to CC Young, February 18, 1933, Box 14, Michigan Department of Health.

[Back](#)

- v. Michigan Department of Health, Whooping Cough Prevention Study, October 13, 1936, Box 4, Kendrick papers.

[Back](#)

- vi. James Doull to Pearl Kendrick, April 25, 1936. See also Doull to Kendrick, March 2, 1936; Kendrick to Doull, May 13, 1936, Kendrick to George Ramsey, September 30, 1936, all Box 3, Kendrick papers.

[Back](#)

- vii. Wade Hampton Frost to Pearl Kendrick, February 2, 1936, Box 3, Kendrick papers.

[Back](#)

- viii. George H. Ramsey to Wade Hampton Frost, August 27, 1936; Kendrick to Frost, February 26, 1936, Box 3, Kendrick papers; Haven Emerson to C.E.A. Winslow, November 18, 1935; Series 1, Box 9, Winslow papers.

[Back](#)

- ix. Wade Hampton Frost to George H. Ramsey, September 10, 1936, Box 3, Kendrick papers.

[Back](#)

- x. Wade Hampton Frost to Pearl Kendrick, August 10, 1936, October 24, 1936, September 21, 1937; Suggestions for Revising Records and Procedures, dictated by Dr. Frost, November 12, 1936; Whooping Cough Immunization Study. Directions for Sorting and Revising Records, September 15, 1937 pursuant to advice of Wade Frost; [WHF], Memorandum for Dr. Kendrick, September 18, 1937; Frost to Kendrick, September 21, 1937 [Boxes 3 and 4, Kendrick papers].

[Back](#)

- xi. Edwin B. Wilson to George M. Lawson, November 19, 1937 [Box 3, Kendrick papers]. Lawson, who had written a Harvard thesis on pertussis, strongly supported Kendrick and had written Wilson for his advice: see George Lawson to Edwin B. Wilson, October 27, 1937. For further discussion of the adequacy of the controls, see Lawson to Wilson, November 22, 1937 and Wilson to Lawson, November 26, 1937. All in Box 2, Wilson papers.]

[Back](#)

- xii. Wade Hampton Frost to Grace Eldering, November 17, 1936, September 21, 1937; Pearl Kendrick to Wade Hampton Frost, October 7, 1936, November 24, 1936; Eldering to Frost, November 25, 1936. All Box 3, Kendrick papers.

[Back](#)

- xiii. Wade Hampton Frost to Haven Emerson, September 12, 1936, Box 4, Kendrick papers.

[Back](#)

- xiv. Frost remained anxious about having a medically qualified epidemiologist--Forsbeck--supervise the data analysis. See George H Ramsey to C.C. Young, April 13, 1937; Young to Ramsey, April 15, 1937. Box 3, Kendrick papers.

[Back](#)

xv. Code Manual, Definitions and Explanations. A Study in Active Immunization Against Pertussis. Michigan Department of Health, Bureau of Laboratories, Western Michigan Division [undated]; Pearl Kendrick to Clark Tibbetts, November 3, 1937. Box 14, Michigan Department of Public Health.

[Back](#)

xvi. Pearl Kendrick to C.C. Young, September 15, 1938, Box 14, Michigan Department of Public Health.

[Back](#)

xvii. Report of the Subcommittee on Evaluation of the Committee on Administrative Practices, [1938], Series III, Box 42, Winslow papers; George Lawson to Pearl Kendrick, November 22, 1937; Pearl Kendrick to James Doull, September 13, 1938, Box 4; Haven Emerson to Pearl Kendrick, March 12, 1938, Box 6, Kendrick papers.

[Back](#)

xviii. See also her comments re EB Wilson's somewhat skeptical letter: Pearl Kendrick to George Lawson, November 30, 1937. Box 4, Kendrick papers.

[Back](#)

[Key Passages](#) [Commentary](#) [Context](#)

[Back](#)

[Home](#)

[Contents](#)